

## Type 2 diabetes

### Definition

Type 2 diabetes is a lifelong (chronic) disease in which there is a high level of sugar (glucose) in the blood. Type 2 diabetes is the most common form of diabetes.

### Causes

Insulin is a hormone produced in the pancreas by special cells, called beta cells. The pancreas is below and behind the stomach. Insulin is needed to move blood sugar (glucose) into cells. Inside the cells, glucose is stored and later used for energy.

When you have type 2 diabetes, your fat, liver, and muscle cells do not respond correctly to insulin. This is called insulin resistance. As a result, blood sugar does not get into these cells to be stored for energy.

When sugar cannot enter cells, a high level of sugar builds up in the blood. This is called hyperglycemia. The body is unable to use the glucose for energy. This leads to the symptoms of type 2 diabetes.

Type 2 diabetes usually develops slowly over time. Most people with the disease are overweight or obese when they are diagnosed. Increased fat makes it harder for your body to use insulin the correct way.

Type 2 diabetes can also develop in people who are thin. This is more common in older adults.

Family history and genes play a role in type 2 diabetes. Low activity level, poor diet, and excess body weight around the waist increase your chance of getting the disease.

### Symptoms


People with type 2 diabetes often have no symptoms at first. They may not have symptoms for many years.

Early symptoms of diabetes caused by a high blood sugar level may include:

- Bladder, kidney, skin, or other infections that are more frequent or heal slowly
- Fatigue
- Hunger
- Increased thirst

### Before


- Include 5 minute warm up
- Check blood sugar



### After

- Include 5 minute cool down
- Check blood sugar

ADAM.



- I.D. bracelet
- Sugar source (glucose tablets or raisins)

ADAM.

- Increased urination
- Blurred vision

## Starchy foods

After many years, diabetes can lead to serious health problems, and as a result, many other symptoms.

### Exams and Tests


Your doctor may suspect that you have diabetes if your blood sugar level is higher than 200 milligrams per deciliter (mg/dL) or 11.1 mmol/L. To confirm the diagnosis, one or more of the following tests must be done.



ADAM.

- Fasting blood glucose level. Diabetes is diagnosed if it is higher than 126 mg/dL (7.0 mmol/L) two different times.
- Hemoglobin A1c (A1C) test. Diabetes is diagnosed if the test result is 6.5% or higher.
- Oral glucose tolerance test. Diabetes is diagnosed if the glucose level is higher than 200 mg/dL (11.1 mmol/L) 2 hours after drinking a special sugar drink.

Low blood sugar symptoms include:



Diabetes screening is recommended for:

- Overweight children who have other risk factors for diabetes, starting at age 10 and repeated every 2 years
- Overweight adults (BMI of 25 or higher) who have other risk factors
- Adults starting at age 45 every 3 years, or at a younger age if the person has risk factors

If you have been diagnosed with type 2 diabetes, you need to work closely with your doctor. See your doctor as often as instructed. This may be every 3 months.

The following exams and tests will help you and your doctor monitor your diabetes and prevent problems.

- Check the skin and bones on your feet and legs.
- Check if your feet are getting numb (diabetic nerve disease).
- Have your blood pressure checked at least once a year (blood pressure goal should be 140/80 mm Hg or lower).
- Have your A1C tested every 6 months if your diabetes is well controlled. Have the test every 3 months if your diabetes is not well controlled.
- Have your cholesterol and triglyceride levels checked once a year.
- Get tests once a year to make sure your kidneys are working well (microalbuminuria and serum creatinine).
- Visit your eye doctor at least once a year, or more often if you have signs of diabetic eye disease.

- See the dentist every 6 months for a thorough dental cleaning and exam. Make sure your dentist and hygienist know that you have diabetes.

## Treat low blood sugar: 15:15 rule



Check blood sugar



Eat 15 grams of carbohydrate



Wait 15 minutes for sugar to get into blood

## Treatment

The goal of treatment at first is to lower your high blood glucose level. Long-term goals are to prevent complications. These are health problems from diabetes.

The most important way to treat and manage type 2 diabetes is by being active and eating healthy foods.

Everyone with diabetes should receive proper education and support about the best ways to manage their diabetes. Ask your doctor about seeing a diabetes nurse educator and a dietitian.


ADAM

## LEARN THESE SKILLS

Learning diabetes management skills will help you live well with diabetes. These skills help prevent health problems and the need for medical care. Skills include:

- How to test and record your blood glucose
- What, when, and how much to eat
- How to safely increase your activity and control your weight
- How to take medicines, if needed
- How to recognize and treat low and high blood sugar
- How to handle sick days
- Where to buy diabetes supplies and how to store them

Your goal is to maintain normal blood glucose levels



ADAM

It may take several months to learn these skills. Keep learning about diabetes, its complications, and how to control and live well with the disease. Stay up-to-date on new research and treatments.

## MANAGING YOUR BLOOD SUGAR

Checking your blood sugar level yourself and writing down the results tells you how well you are managing your diabetes. Talk to your doctor and diabetes educator about how often to check.

To check your blood sugar level, you use a device called a glucose meter. Usually, you prick your finger with a small needle called a lancet. This gives you a tiny drop of blood. You place the blood on a test strip and put the strip into the meter. The meter gives you a reading that tells you the level of your blood sugar.

Your doctor or diabetes educator will help set up a testing schedule for you. Your doctor will help you set a target range for your blood sugar numbers. Keep these factors in mind:

- Most people with type 2 diabetes only need to check their blood sugar once or twice a day.
- If your blood sugar level is under control, you may only need to check it a few times a week.
- You may test yourself when you wake up, before meals, and at bedtime.

- You may need to test more often when you are sick or under stress.
- You may need to test more often if you are having more frequent low blood sugar symptoms.

Keep a record of your blood sugar for yourself and your doctor. Based on your numbers, you may need to make changes to your meals, activity, or medicines to keep your blood sugar level in the right range.

#### HEALTHY EATING AND WEIGHT CONTROL

Work closely with your health care providers to learn how much fat, protein, and carbohydrates you need in your diet. Your meal plans should fit your lifestyle and habits and should include foods that you like.

Managing your weight and having a well-balanced diet are important. Some people with type 2 diabetes can stop taking medicines after losing weight. This does not mean that their diabetes is cured. They still have diabetes.

Very obese people whose diabetes is not well managed with diet and medicine may consider weight loss (bariatric) surgery.

#### REGULAR PHYSICAL ACTIVITY

Regular activity is important for everyone. It is even more important when you have diabetes. Exercise is good for your health because it:

- Lowers your blood sugar level without medicine
- Burns extra calories and fat to help manage your weight
- Improves blood flow and blood pressure
- Increases your energy level
- Improves your ability to handle stress


Talk to your doctor before starting any exercise program. People with type 2 diabetes may need to take special steps before, during, and after physical activity or exercise.

#### MEDICINES TO TREAT DIABETES

If diet and exercise do not help keep your blood sugar at normal or near-normal levels, your doctor may prescribe medicine. Since these drugs help lower your blood sugar level in different ways, your doctor may have you take more than one drug.

Some of the most common types of medicines are listed below. They are taken by mouth or injection.


- Alpha-glucosidase inhibitors
- Biguanides
- DPP IV inhibitors



Alpha-glucosidase inhibitors slow the digestion of starches



ADAM




Biguanides stop the liver from making extra sugar when it's not needed



ADAM

- Injectable medicines (GLP-1 analogs)
- Meglitinides
- SGL T2 inhibitors
- Sulfonylureas
- Thiazolidinediones

You may need to take insulin if your blood sugar cannot be controlled with any of the above medicines. Most commonly, insulin is injected under the skin using a syringe, insulin pen, or pump. Another form of insulin is the inhaled type. Insulin cannot be taken by mouth because the acid in the stomach destroys the insulin.



Sulfonylureas help your pancreas make extra insulin




ADAM.

## PREVENTING COMPLICATIONS

Your doctor may prescribe medicines or other treatments to reduce your chance of developing some of the more common complications of diabetes, including:

- Eye disease
- Kidney disease
- Heart disease and stroke



Thiazolidinediones help body cells use insulin better



ADAM.

## FOOT CARE

People with diabetes are more likely than those without diabetes to have foot problems. Diabetes damages the nerves. This can make your feet less able to feel pressure, pain, heat, or cold. You may not notice a foot injury until you have severe damage to the skin and tissue below, or you get a severe infection.

Diabetes can also damage blood vessels. Small sores or breaks in the skin may become deeper skin sores (ulcers). The affected limb may need to be amputated if these skin ulcers do not heal or become larger, deeper, or infected.

To prevent problems with your feet:

- Stop smoking if you smoke.
- Improve control of your blood sugar.
- Get a foot exam by your doctor at least twice a year to learn if you have nerve damage.
- Check and care for your feet every day. This is very important when you already have nerve or blood vessel damage or foot problems.
- Treat minor infections, such as athlete's foot, right away.
- Use moisturizing lotion on dry skin.
- Make sure you wear the right kind of shoes. Ask your doctor what type of shoe is right for you.

## Support Groups

There are many diabetes resources that can help you understand more about type 2 diabetes. You can also learn ways to manage your condition so you can live well with diabetes.

## Outlook (Prognosis)


Diabetes is a lifelong disease and there is no cure.

Some people with type 2 diabetes no longer need medicine if they lose weight and become more active. When they reach their ideal weight, their body's own insulin and a healthy diet can control their blood sugar level.


## Possible Complications

After many years, diabetes can lead to serious health problems:

- You could have eye problems, including trouble seeing (especially at night), and light sensitivity. You could become blind.
- Your feet and skin can develop sores and infections. After a long time, your foot or leg may need to be amputated. Infection can also cause pain and itching in other parts of the body.
- Diabetes may make it harder to control your blood pressure and cholesterol. This can lead to a heart attack, stroke, and other problems. It can become harder for blood to flow to your legs and feet.
- Nerves in your body can get damaged, causing pain, tingling, and numbness.
- Because of nerve damage, you could have problems digesting the food you eat. You could feel weakness or have trouble going to the bathroom. Nerve damage can make it harder for men to have an erection.
- High blood sugar and other problems can lead to kidney damage. Your kidneys may not work as well as they used to. They may even stop working so that you need dialysis or a kidney transplant.



ADAM.



## When to Contact a Medical Professional

Call 911 right away if you have:

- Chest pain or pressure
- Fainting, confusion or unconsciousness
- Seizure
- Shortness of breath

These symptoms can quickly get worse and become emergency conditions (such as convulsions, hypoglycemic coma or hyperglycemic coma).

Also call your doctor if you have:

- Numbness, tingling, or pain in your feet or legs
- Problems with your eyesight
- Sores or infections on your feet
- Symptoms of high blood sugar (extreme thirst, blurry vision, dry skin, weakness or fatigue, the need to urinate a lot)
- Symptoms of low blood sugar (weakness or fatigue, trembling, sweating, irritability, trouble thinking clearly, fast heartbeat, double or blurry vision, uneasy feeling)

## Prevention

You can help prevent type 2 diabetes by staying at a healthy body weight. You can get to a healthy weight by eating healthy foods, controlling your portion sizes, and leading an active lifestyle. Some medicines can also delay or prevent type 2 diabetes in people at risk of developing the disease.

## Alternative Names

Noninsulin-dependent diabetes; Diabetes - type II; Adult-onset diabetes; Diabetic - type 2 diabetes; Oral hypoglycemic - type 2 diabetes

---

Review Date: July 24, 2015.

Reviewed By: Brent Wisse, MD, Associate Professor of Medicine, Division of Metabolism, Endocrinology & Nutrition, University of Washington School of Medicine, Seattle, WA. Internal review and update on 09/01/2016 by David Zieve, MD, MHA, Isla Ogilvie, PhD, and the A.D.A.M. Editorial team.



A.D.A.M., Inc. is accredited by URAC, also known as the American Accreditation HealthCare Commission ([www.urac.org](http://www.urac.org)). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#) and [privacy policy](#). A.D.A.M. is also a founding member of Hi-Ethics and subscribes to the principles of the Health on the Net Foundation ([www.hon.ch](http://www.hon.ch)).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Call 911 for all medical emergencies. Links to other sites are provided for information only -- they do not constitute endorsements of those other sites. © 1997- 2017 A.D.A.M., Inc. Any duplication or distribution of the information contained herein is strictly prohibited.

